

N-flat connections

Mauricio Angel and Rafael Díaz

February 2, 2008

Abstract

We construct geometric examples of N -differential graded algebras such as the algebra of differential forms of depth N on an affine manifold, and N -flat covariant derivatives.

Introduction

The applications of the theory of complexes and homological algebra have touched many branches of mathematics such as topology, geometry and mathematical physics. The possibility of developing an homological algebra for the equation $d^N = 0$ for $N \geq 3$ has been around at least since 1940 in works by Mayer [M].

However, N -homological algebra for $N \geq 3$ only aroused the interest that it deserves in 1991 when Kapranov's paper [K] appeared. Fundamental works by Dubois-Violette appeared in [DV1], [DV2] soon after. The key notion is that of a q -differential graded algebra (q must be a primitive N -root of unity), which consists of a $\mathbb{Z}$ -graded vector space V together with an operation $m : V \otimes V \rightarrow V$ and an degree one map $d : V \rightarrow V$ such that: 1) m is associative, 2) d satisfy the q -Leibniz rule $d(ab) = d(a)b + q^{\deg(a)}ad(b)$ and 3) $d^N = 0$.

Notice that in the definition of a q -differential graded algebra not only the equation $d^2 = 0$ is deformed to $d^N = 0$, but also the Leibniz rule is substituted by the q -Leibniz rule. The theory of q -differential graded algebras has been further developed in many papers such as [AB], [DV3], [DVK], [KW], [KN], [S].

The foundations of a theory of N -differential graded algebras (N -dga) has been written down in [AD]. A N -dga consists of a $\mathbb{Z}$ -graded vector space V together with an associative operation $m : V \otimes V \rightarrow V$ and a degree one map $d : V \rightarrow V$ such that: 1) $d^N = 0$ and 2) $d(ab) = d(a)b + (-1)^{\deg(a)}ad(b)$. Notice that in the definition of a N -dga the equation $d^2 = 0$ is replaced by $d^N = 0$, whereas the Leibniz rule is not modified. This fact explain why our definition is better suited for the use of differential geometric techniques.

At first sight it looks difficult to come up with examples of N-dga. The purpose of this paper is to introduce geometric examples of N-differential graded algebras, which show that that this sort of algebras appear naturally in a wide variety of contexts. Our examples range from the algebra of differential forms of depth N (see Section 4) to the theory of N-flat covariant derivatives (see Section 1).

1 N-covariant differentials

The purpose of this paper is to introduce geometric examples of N-differential graded algebras (N-dga). Let us first formally introduce the notion of N-dga [AD]. Let $\mathbf{k}$ be a commutative ring with unit.

Definition 1 *A N-differential graded algebra or N-dga over $\mathbf{k}$, is a triple $(A^\bullet, m, d)$ where $m : A^k \otimes A^l \rightarrow A^{k+l}$ and $d : A^k \rightarrow A^{k+1}$ are $\mathbf{k}$ -modules homomorphisms satisfying*

- 1) $(A^\bullet, m)$ is a graded associative algebra.
- 2) d satisfies the graded Leibniz rule $d(ab) = d(a)b + (-1)^{\bar{a}}ad(b)$.
- 3) $d^N = 0$, i.e., $(A^\bullet, d)$ is a N-complex.

A 1-dga is a graded associative algebra. A 2-dga is a differential graded algebra.

Definition above may be justified categorically as follows: consider the category $\mathbf{NComp}_{\mathbf{k}}$ of nilpotent differential graded $\mathbf{k}$ -modules. Objects in $\mathbf{NComp}_{\mathbf{k}}$ are pairs (V, d) where V is a $\mathbb{Z}$ -graded $\mathbf{k}$ -module $V = \bigoplus_{i \in \mathbb{Z}} V^i$ together with a degree one map $d : V \rightarrow V$ such that $d^N = 0$ for some $N \geq 2$. $\mathbf{NComp}_{\mathbf{k}}$ is a symmetric monoidal category since if (V_1, d_1) is such that $d_1^{N_1} = 0$ and (V_2, d_2) is such that $d_2^{N_2} = 0$ then $(V_1 \otimes V_2, d_1 \otimes Id + Id \otimes d_2)$ satisfy $(d_1 \otimes Id + Id \otimes d_2)^{N_1+N_2-1} = 0$. N-differential graded algebras (for $N \geq 2$) are the monoids in $\mathbf{NComp}_{\mathbf{k}}$.

Definition 2 *A N-dga $(A^\bullet, m, d)$ is called **proper** if $d^{N-1} \neq 0$.*

Let M be a smooth finite dimensional manifold and $\mathcal{E} = M \times E$ be a trivial bundle over M . The reader may assume that $\mathcal{E}$ is a global bundle and that M is actually a neighborhood on which $\mathcal{E}$ is trivial. Since our results are covariant they will hold globally as well.

The space $\Omega^\bullet(M, End(\mathcal{E}))$ of $End(\mathcal{E})$ -valued forms on M is endowed with a differential graded algebra structure, with the product given by

$$\begin{aligned} \wedge : \Omega^\bullet(M, End(\mathcal{E})) \otimes \Omega^\bullet(M, End(\mathcal{E})) &\rightarrow \Omega^\bullet(M, End(\mathcal{E})) \\ (\alpha \otimes \psi)(\beta \otimes \phi) &\longmapsto (\alpha \wedge \beta) \otimes \psi\phi. \end{aligned}$$

The space of $\mathcal{E}$ -valued forms $\Omega^\bullet(M, \mathcal{E})$ is endowed with a differential graded module structure over $\Omega^\bullet(M, \text{End}(\mathcal{E}))$ and also over the differential graded algebra $\Omega^\bullet(M)$ of differential forms on M , given by

$$\begin{aligned}\Omega^\bullet(M) \otimes \Omega^\bullet(M, \mathcal{E}) &\rightarrow \Omega^\bullet(M, \mathcal{E}) \\ \alpha(\beta \otimes \phi) &\mapsto (\alpha \wedge \beta) \otimes \phi.\end{aligned}$$

$$\begin{aligned}\Omega^\bullet(M, \text{End}(\mathcal{E})) \otimes \Omega^\bullet(M, \mathcal{E}) &\rightarrow \Omega^\bullet(M, \mathcal{E}) \\ (\alpha \otimes \Phi)(\beta \otimes \phi) &\mapsto (\alpha \wedge \beta) \otimes \Phi(\phi)\end{aligned}$$

Recall that a covariant derivative ∇ on $\mathcal{E}$ is a linear map $\nabla : \Omega^\bullet(M, E) \rightarrow \Omega^\bullet(M, E)$ of degree 1 such that

$$\nabla(a\alpha) = (da)\alpha + (-1)^{\deg(a)}a\nabla\alpha, \text{ for all } a \in \Omega^\bullet(M), \alpha \in \Omega^\bullet(M, E).$$

Since $\mathcal{E}$ is trivial, ∇ may be written as $\nabla = d + \omega$ for some $\omega \in \Omega^1(M, \text{End}(\mathcal{E}))$, where d is the de Rham differential and ω is the connection one-form of ∇ .

For any $\alpha \in \Omega^k(M, E)$, $\nabla\alpha \in \Omega^{k+1}(M, E)$ is the E -valued form given by

$$\nabla\alpha = d\alpha + \omega \wedge \alpha. \quad (1.1)$$

Taking the covariant derivative of (1.1) we get

$$\nabla^2(\alpha) = d\omega \wedge \alpha + \omega \wedge \omega \wedge \alpha = (d\omega + \omega \wedge \omega) \wedge \alpha = F_\omega \wedge \alpha. \quad (1.2)$$

The 2-form $F_\omega \in \Omega^2(M, \text{End}(\mathcal{E}))$ is called the curvature of ∇ . A connection ω is said to be **flat** if $\nabla^2 = 0$ or equivalently if $F_\omega = 0$.

Definition 3 We say that a connection ω is **N-flat** if $\nabla^N = 0$, where $\nabla = d + \omega$ and $N \geq 2$.

A 2-flat connection is just a flat connection. We have the following

Theorem 4 Let M be a manifold and $\mathcal{E} = M \times E$ be a trivial bundle over M . A connection ω is

1. $2N$ -flat if and only if $F_\omega^N = 0$.
2. $(2N + 1)$ -flat if and only if $F_\omega^N \nabla = 0$.

proof Suppose that $K = 2N + n$, $n \in \{0, 1\}$, then

$$\nabla^K = (d + \omega)^{2N+n} = (d(\omega) + \omega \wedge \omega)^N \nabla^n = F_\omega^N \nabla^n. \blacklozenge$$

Example 5 Consider $\mathbb{R}^4$ with coordinates (x_1, x_2, x_3, x_4) . For $\omega_1, \omega_2 : \mathbb{R}^4 \rightarrow \mathbb{R}$ we consider the connection $\omega = \omega_1 dx_1 + \omega_2 dx_2$.

A simple calculation shows that

$$F_\omega = \left(\frac{\partial \omega_1}{\partial x_2} - \frac{\partial \omega_2}{\partial x_1} \right) dx_1 \wedge dx_2 \neq 0, \text{ if } \frac{\partial \omega_1}{\partial x_2} - \frac{\partial \omega_2}{\partial x_1} \neq 0.$$

For example one can take $\omega_1 = x_2$ and $\omega_2 = -x_1$, then $F_\omega = 2dx_1 \wedge dx_2 \neq 0$. But $F_\omega^2 = 0$ which implies that ω is a 4-flat connection and $(\Omega^\bullet(\mathbb{R}^4, \text{End}(\mathbb{R}^4)), \nabla)$ is a proper 4-dga.

The previous example is an instance of the following result.

Theorem 6 Let M be a manifold and assume that $TM = A \oplus B$, where $\dim(A) = a$, $\dim(B) = b$, and ω is a connection on TM such that if $F_\omega(\alpha, \beta) \neq 0$ then $\alpha \in A$ and $\beta \in A$. In this case $\nabla = d + \omega$ is $2N$ -flat for $N > a$.

Let now M be a n -dimensional smooth manifold and $\mathcal{E} = M \times E$ a trivial bundle on it. Using local coordinates $x_1, \dots, x_n$ a connection $\omega \in \Omega^1(M, \text{End}(\mathcal{E}))$ may be written as $\omega = \omega_i dx^i$. The 2-form $d\omega + \omega \wedge \omega$ can be written as $F_{ij} dx^i \wedge dx^j$ where $F_{ij} = \partial_i \omega_j - \partial_j \omega_i + [\omega_i, \omega_j]$. Furthermore

$$(F_{ij} dx^i \wedge dx^j)^k = \sum_{A \subseteq [n]} \left(\sum_{\alpha \in P(A)} \prod_{i=1}^k \text{sign}(\alpha) F_{a_i, b_i} \right) dx_{s_1} \wedge \dots \wedge dx_{s_{2k}}$$

where $[n]$ denotes the set $\{1, 2, \dots, n\}$, the cardinality of A is $2k$ and $P(A)$ is the set of ordered pairings of $A = \{s_1 < \dots < s_{2k}\}$. An ordered pairing $\alpha \in P(A)$ is a sequence $\{(a_i, b_i)\}_{i=1}^k$ such that $A = \bigsqcup_{i=1}^k \{a_i, b_i\}$ and $a_i < b_i$. Theorem 4 part (1) implies

Theorem 7 $(\Omega^\bullet(M, \mathcal{E}), d + \omega)$ is a $2k$ -dga if and only if

$$\sum_{\alpha \in P(A)} \text{sign}(\alpha) \prod_{i=1}^k F_{a_i, b_i} = 0,$$

for all $A \subseteq [n]$ with cardinality $2k$.

Corollary 8 If $\dim(M) = 2n$, then $(\Omega^\bullet(M, \mathcal{E}), d + \omega)$ is a $2n$ -dga if and only if

$$\sum_{\alpha \in P([2n])} \text{sign}(\alpha) \prod_{i=1}^n F_{a_i, b_i} = 0.$$

Let (M, g) be a Riemannian manifold. The tangent bundle TM has a canonical covariant derivative ∇ , called the Levi-Civita connection. Suppose that the Riemannian metric is given in local coordinates $x_1, \dots, x_n$ by the positive definite symmetric matrix g_{ij} :

$$ds^2 = \sum g_{ij} dx^i dx^j.$$

Denote by $\partial_1, \dots, \partial_n$ the corresponding coordinate vector fields, then the covariant derivatives can be expressed

$$\nabla_{\partial_i} \partial_j = \sum_k \Gamma_{jk}^i \partial_k \quad \Gamma_{jk}^i = \sum_l \frac{1}{2} (\partial_k g_{lj} + \partial_j g_{lk} - \partial_l g_{jk}) g^{il},$$

where (g^{lk}) is the inverse matrix to (g_{kl}) . In this case the curvature 2-form is given by

$$R = R_{kl} dx_k \wedge dx_l = R_{jkl}^i \partial_j \otimes dx_i \otimes dx_k \wedge dx_l,$$

where

$$R_{jkl}^i = \partial_k \Gamma_{jl}^i - \partial_l \Gamma_{jk}^i + \Gamma_{jl}^h \Gamma_{hk}^i - \Gamma_{jk}^h \Gamma_{hl}^i.$$

Theorem 9 $(\Omega^\bullet(M, TM), \nabla)$ is a $2k$ -dga if and only if

$$\sum_{\alpha \in P(A)} \text{sign}(\alpha) \prod_{i=1}^k R_{a_i b_i k}^l = 0,$$

for all $A \subseteq [n]$ with cardinality $2k$.

Example 10 Consider the space $S^2 \times T^2$ with coordinates (x_1, x_2, x_3, x_4) , $0 < x_1 < 2\pi$, $-\pi/2 < x_2 < \pi/2$, $0 < x_3 < 1$ and $0 < x_4 < 1$. The metric on $S^2 \times T^2$ is given in those coordinates by

$$ds^2 = \sin^2(x_2)(dx_1)^2 + (dx_2)^2 + (dx_3)^2 + (dx_4)^2.$$

The curvature is

$$R = \begin{pmatrix} 0 & 1 \\ -\sin^2(x_2) & 0 \end{pmatrix} dx_1 \wedge dx_2 \neq 0$$

Since $R^2 = 0$, then $(\Omega(S^2 \times \mathbb{R}^2, T(S^2 \times T^2)), \nabla)$ is a 4-dga.

We generalize the previous example as follows:

Theorem 11 Let (M, g) be a Riemannian manifold and assume that $TM = A \oplus B$ is an orthogonal decomposition, where $\dim(A) = a$, $\dim(B) = b$, and g is flat on B , then $(\Omega(M, TM), \nabla)$ is a $2N$ -dga for $N \geq a$.

Recall the definition of the generalized cohomology groups of a N-complex (A, d) [Kap]

$${}_p H^i(A) = \frac{\text{Ker}\{d^p : A^i \rightarrow A^{i+p}\}}{\text{Im}\{d^{N-p} : A^{i-N+p} \rightarrow A^i\}}.$$

The total object $\mathbf{H}^\bullet(A)$ of the cohomology associated to a N -complex is $\mathbf{H}^\bullet(A) = \bigoplus_{m=0}^\infty \mathbf{H}^m(A)$, where

$$\mathbf{H}^m(A) = \bigoplus_{2i-p=m} {}_p H^i(A).$$

Example 12 Consider the torus T^4 with global coordinates $(\theta_1, \theta_2, \theta_3, \theta_4)$ satisfying $(\theta_1, \theta_2, \theta_3, \theta_4) = (\theta_1 + 1, \theta_2 + 1, \theta_3 + 1, \theta_4 + 1)$. For

$$E_{11} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad E_{12} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

We define $\nabla = d + E_{11}d\theta_1 + E_{12}d\theta_2$, then since $[E_{11}, E_{12}] = E_{12}$ we have $\nabla^2 = E_{12}d\theta_1 \wedge d\theta_2 \neq 0$, nevertheless it is obvious that $\nabla^4 = 0$ thus $(\Omega(T^4, \mathbb{C}^2), \nabla)$ is a (proper) 4-dga.

We have $(d\theta_1 \wedge d\theta_2)(d\theta_1) = 0$ which implies that $d\theta_1 \in \text{Ker}(\nabla^2)$, and $d\theta_1$ is not in $\text{Im}(\nabla^2)$, because otherwise $\nabla^2\beta$ is a 2-form for all functions β . Then the cohomology group ${}_2 H^1(T^4) \neq 0$.

In [AD] we showed that if (A, d) is a N -dga then $(\mathbf{H}(A), d)$ is a $(N-1)$ -dga, thus we see that $\mathbf{H}(\Omega(T^4, \mathbb{C}^2), \nabla) (\neq 0)$ is a (proper) 3-dga.

Let M be a manifold and $\mathcal{E}_i$ trivial bundles on M , $i = 1, 2$. For $\nabla^{\mathcal{E}_i}$ covariant derivatives on $\mathcal{E}_i$, we have a natural covariant derivative $\nabla^{\mathcal{E}_1 \otimes \mathcal{E}_2}$ on the tensor product $\mathcal{E}_1 \otimes \mathcal{E}_2$ given by

$$\nabla^{\mathcal{E}_1 \otimes \mathcal{E}_2}(s_1 \otimes s_2) = \nabla^{\mathcal{E}_1} s_1 \otimes s_2 + s_1 \otimes \nabla^{\mathcal{E}_2} s_2,$$

for $s_i \in \Omega(M, \mathcal{E}_i)$.

The next result is an analog to Theorem 8 [AD] and we omit the proof,

Theorem 13 If $(\Omega(M, \mathcal{E}_1), \nabla^{\mathcal{E}_1})$ is a N -dga and $(\Omega(M, \mathcal{E}_2), \nabla^{\mathcal{E}_2})$ is a M -dga, then $(\Omega(M, \mathcal{E}_1 \otimes \mathcal{E}_2), \nabla^{\mathcal{E}_1 \otimes \mathcal{E}_2})$ is a $(M+N-1)$ -dga. Equivalently, if $\nabla^{\mathcal{E}_1}$ is M -flat and if $\nabla^{\mathcal{E}_2}$ is N -flat then $\nabla^{\mathcal{E}_1 \otimes \mathcal{E}_2}$ is $(M+N-1)$ -flat.

2 N-Chern-Simons actions

Let M be a manifold, $\dim(M) = 2K+1$ and $\mathcal{E} = M \times E$ be a trivial bundle over M . For a connection ω on $\mathcal{E}$, consider its K-Curvature $F_\omega^K = (d\omega + \omega \wedge \omega)^K$. On $\Omega^\bullet(M, E)$

exists a linear functional $\int_M Tr: \Omega^\bullet(M, End(E)) \rightarrow \mathbb{R}$ of degree $2K+1$, i.e., $\int b = 0$ if $\bar{b} \neq 2K+1$, given by

$$\omega \longmapsto \int_M Tr(\omega), \text{ for all } \omega \in \Omega^\bullet(M, End(E)).$$

The functional $\int_M Tr$ satisfy the conditions of [AD],

1. $\int_M Tr$ is non degenerate, that is, $\int_M Tr(\alpha \wedge \beta) = 0$ for all α , then $\beta = 0$.
2. $\int_M Tr(d(\alpha)) = 0$ for all α , where $d = d_{End(E)}$.
3. $\int_M Tr$ is cyclic, this is

$$\int_M Tr(\alpha_1 \alpha_2 \dots \alpha_n) = (-1)^{\bar{\alpha}_1(\bar{\alpha}_2 + \dots + \bar{\alpha}_n)} \int_M Tr(\alpha_2 \dots \alpha_n \alpha_1).$$

We define the **Chern-Simons** functional $cs_{2,2K}: \Omega^\bullet(M, End(E)) \rightarrow \mathbb{R}$ by

$$cs_{2,2K}(\omega) = 2K \int_M Tr(\pi(\#^{-1}(\omega(d\omega + \omega^2)^K)))$$

where

1. $\mathbb{R}\langle\omega, d\omega\rangle$ denotes the free $\mathbb{R}$ -algebra generated by symbols ω and $d\omega$.
2. $\#: \mathbb{R}\langle\omega, d\omega\rangle \longrightarrow \mathbb{R}\langle\omega, d\omega\rangle$ is the linear map defined by

$$\#(\omega^{i_1} d(\omega)^{j_1} \dots \omega^{i_k} d(\omega)^{j_k}) = (i_1 + \dots + i_k + j_1 + \dots + j_k) \omega^{i_1} d(\omega)^{j_1} \dots \omega^{i_k} d(\omega)^{j_k}.$$

3. $\pi: \mathbb{R}\langle\omega, d(\omega)\rangle \longrightarrow \Omega^\bullet(M, End(E))$ is the canonical projection.

We have the following result

Theorem 14 *Let $K \geq 1$ be an integer. The Chern-Simons functional $cs_{2,2K}$ is the Lagrangian for the $2K$ -Maurer-Cartan equation, i.e., $\omega \in \Omega^\bullet(M, End(E))$ is a critical point of $cs_{2,2K}$ if and only if $F_\omega^K = 0$.*

For $K = 2, 3, 4$ the Chern-Simons functional $cs_{2,2K}(a)$ is given by

$$\begin{aligned} cs_{2,4}(\omega) &= \int_M Tr\left(\frac{4}{3}\omega(d(\omega))^2 + 2\omega^3 d(\omega) + \frac{4}{5}\omega^5\right). \\ cs_{2,6}(\omega) &= \int_M Tr\left(\frac{3}{2}\omega(d(\omega))^3 + \frac{18}{5}\omega^3(d(\omega))^2 + 3\omega^5 d(\omega) + \frac{6}{7}\omega^7\right). \\ cs_{2,8}(\omega) &= \int_M Tr\left(\frac{8}{5}\omega(d(\omega))^4 + \frac{16}{3}\omega^3(d(\omega))^3 + \frac{48}{7}\omega^5(d(\omega))^2 + 4\omega^7 d(\omega) + \frac{8}{9}\omega^9\right). \end{aligned}$$

3 (K,N)-flat connections

Suppose now that we choose a nilpotent derivation $\delta \in \Omega^1(M, \text{End}(\mathcal{E}))$ instead of the exterior differential, in this case any covariant derivative on $M \times E$ still may be written $\nabla = \delta + \omega$ for some $\omega \in \Omega^1(M, \text{End}(\mathcal{E}))$. Before continue we will review some notations from [AD].

For $s = (s_1, \dots, s_n) \in \mathbb{N}^n$ we set $l(s) = n$, the length of the vector s , and $|s| = \sum_i s_i$. For $1 \leq i < n$, $s_{>i}$ denotes the vector given by $s_{>i} = (s_{i+1}, \dots, s_n)$, for $1 < i \leq n$, $s_{<i}$ stands for $s_{<i} = (s_1, \dots, s_{i-1})$, we also set $s_{>n} = s_{<1} = \emptyset$. $\mathbb{N}^{(\infty)}$ denotes the set $\bigsqcup_{n=0}^{\infty} \mathbb{N}^n$, where by convention $\mathbb{N}^0 = \{\emptyset\}$.

For $e \in \Omega^1(M, \text{End}(\mathcal{E}))$, we define $e^{(s)} = e^{(s_1)} \dots e^{(s_n)}$, where $e^{(a)} = d_{\text{End}}^a(e)$ if $a \geq 1$, $e^{(0)} = e$ and $e^\emptyset = 1$. In the case that $e_\omega \in \Omega^1(M, \text{End}(\mathcal{E}))$ given by

$$e_\omega(\alpha) = \omega \wedge \alpha, \quad \omega \in \Omega^1(M)$$

then $e_\omega^{(a)} = d_{\text{End}}^a(e_\omega)$ reduce to $e_\omega^{(a)} = e_{d^a(\omega)}$, thus

$$e_\omega^{(s)} = e_\omega^{(s_1)} \dots e_\omega^{(s_n)} = e_{d^{s_1}(\omega)} \dots e_{d^{s_n}(\omega)}.$$

For $N \in \mathbb{N}$ we define $E_N = \{s \in \mathbb{N}^{(\infty)} : |s| + l(s) \leq N\}$ and for $s \in E_N$ we define $N(s) \in \mathbb{Z}$ by $N(s) = N - |s| - l(s)$.

We introduce the discrete quantum mechanical system L by

1. $V_L = \mathbb{N}^{(\infty)}$.
2. There is a unique directed edge in L from vertex s to t if and only if $t \in \{(0, s), s, (s + e_i)\}$ where $e_i = (0, \dots, \underbrace{1}_{i\text{-th}}, \dots, 0) \in \mathbb{N}^{l(s)}$.
3. Edges in L are weighted according to the following table

$s(e_i)$	$t(e_i)$	$v(e_i)$
s	$(0, s)$	1
s	s	$(-1)^{ s +l(s)}$
s	$(s + e_i)$	$(-1)^{ s_{<i} +i-1}$

The set $P_N(\emptyset, s)$ consist of all paths $\gamma = (e_1, \dots, e_N)$, such that $s(e_1) = \emptyset$ and $t(e_N) = s$. For $\gamma \in P_N(\emptyset, s)$ we define the weight $v(\gamma)$ of γ as

$$v(\gamma) = \prod_{i=1}^N v(e_i).$$

For example if we have $\delta^K = 0$ then the N -curvature is given by

Theorem 15 *Let M be a manifold and $\mathcal{E} = M \times E$ be a trivial bundle over M . Given any connection ω consider the covariant differential $\nabla = \delta + \omega$, then the N -curvature is given by*

$$\nabla^N = \sum_{k=0}^{N-1} c_k \delta^k$$

where

$$c_k = \sum_{\substack{s \in E_N \\ N(s)=k \\ s_i < K}} c(s, N) \omega^{(s)} \quad \text{and} \quad c(s, N) = \sum_{\gamma \in P_N(\emptyset, s)} v(\gamma).$$

Suppose that we have a derivation δ such that $\delta^3 = 0$, and we want to deform it into a covariant 3-differential $\nabla = \delta + \omega$, $\omega \in \Omega^1(M)$. So we required that $\nabla^3 = 0$. By Theorem 15 we must have $\sum_{k=0}^2 c_k \delta^k = 0$, let us calculate the coefficients c_k . Notice that

$$E_3 = \{\emptyset, (0), (1), (2), (0, 0), (1, 0), (0, 1), (0, 0, 0)\}$$

Let us first compute $\mathbf{c}_0$. In this case we have four vectors in E_3 such that $N(s) = 0$, these are $(2), (1, 0), (0, 1), (0, 0, 0)$.

For $s = (2)$ there is only one path from $\emptyset$ to (2) of length 3,
 $\emptyset \rightarrow (0) \rightarrow (1) \rightarrow (2)$ with weight is 1 and $\omega^{(2)} = \delta^2(\omega)$, thus $c(s, 3) = \delta^2(\omega)$.

For $s = (1, 0)$ there is only one path from $\emptyset$ to $(1, 0)$ of length 3,
 $\emptyset \rightarrow (0) \rightarrow (0, 0) \rightarrow (1, 0)$ with weight 1 and $\omega^{(1,0)} = \delta(\omega)\omega$, thus $c(s, 3) = \delta(\omega)\omega$.

For $s = (0, 1)$ the paths from $\emptyset$ to $(0, 1)$ of length 3 are
 $\emptyset \rightarrow (0) \rightarrow (0, 0) \rightarrow (0, 1)$ with weight -1.
 $\emptyset \rightarrow (0) \rightarrow (1) \rightarrow (0, 1)$ with weight 1.
 $\omega^{(0,1)} = \omega\delta(\omega)$, then $c(s, 3) = 0$, because the sum of the weight of the 2 paths is 0.

For $s = (0, 0, 0)$ the only path from $\emptyset$ to $(0, 0, 0)$ of length 3 is
 $\emptyset \rightarrow (0) \rightarrow (0, 0) \rightarrow (0, 0, 0)$ the weight is 1 and $\omega^{(0,0,0)} = \omega^3$, thus $c(s, 3) = \omega^3$.

We proceed to compute $\mathbf{c}_1$. In this case we have two vectors in E_3 such that $N(s) = 1$, (1) and $(0, 0)$

For $s = (1)$, the paths from $\emptyset$ to (1) of length 3 are
 $\emptyset \rightarrow \emptyset \rightarrow (0) \rightarrow (1)$ with weight 1.
 $\emptyset \rightarrow (0) \rightarrow (0) \rightarrow (1)$ with weight -1.
 $\emptyset \rightarrow (0) \rightarrow (1) \rightarrow (1)$ with weight 1.
 $\omega^{(1)} = \delta(\omega)$ and thus $c(s, 3) = \delta(\omega)$.

For $s = (0, 0)$ the paths from $\emptyset$ to $(0, 0)$ of length 3 are
 $\emptyset \rightarrow (0) \rightarrow (0) \rightarrow (0, 0)$ with weight -1.
 $\emptyset \rightarrow \emptyset \rightarrow (0) \rightarrow (0, 0)$ with weight 1.
 $\emptyset \rightarrow (0) \rightarrow (0, 0) \rightarrow (0, 0)$ with weight 1.
 $\omega^{(0,0)} = \omega^2$ and thus $c(s, 3) = \omega^2$.

Finally we compute $\mathbf{c}_2$. In this case we have one vector in E_3 such that $N(s) = 2, (0)$

For the case $s = (0)$ the paths from $\emptyset$ to (0) of length 3 are
 $\emptyset \rightarrow \emptyset \rightarrow \emptyset \rightarrow (0)$ with weight 1.
 $\emptyset \rightarrow \emptyset \rightarrow (0) \rightarrow (0)$ with weight -1.
 $\emptyset \rightarrow (0) \rightarrow (0) \rightarrow (0)$ with weight 1.
 $\omega^{(0)} = \omega$ and thus $c(s, 3) = \omega$.

Now the 3-curvature is given by

$$\nabla^3 = (\delta^2(\omega) + \delta(\omega)\omega + \omega^3) + (\delta(\omega) + \omega^2)\delta + \omega\delta^2.$$

Example 16 *Infinitesimal deformations.* Suppose that we have a K -differential δ and for a connection ω we consider the infinitesimal deformation $\nabla = \delta + t\omega$, where $t^2 = 0$. ω is N -flat if we have $\sum_k c_k \delta^k = 0$, but in this case

$$(t\omega)^{(s)} = (t\omega)^{(s_1)} \dots (t\omega)^{(s_l(s))} = t^{l(s)} \omega^{(s)} = 0 \quad \text{unless } l(s) \leq 1.$$

Thus E_N is given by

$$E_N = \{(1), \dots, (K-1)\}$$

and $c_k = c((N - (k+1)), N) \omega^{(N-(k+1))}$ for all k such that $2 \leq N - k \leq K$.

4 Differential forms of depth N

We consider $U \subseteq \mathbb{R}^k$ an open set. We use local coordinates $x_1, \dots, x_k$ on U . For $(N_1, \dots, N_k) \in \mathbb{N}_{\geq 2}^k$ we define the algebra $\Omega_{(N_1, \dots, N_k)}(U)$ of differential forms of depth $(N_1, \dots, N_k)$ on $\mathbb{R}^k$ as follows: an element $\alpha \in \Omega_{(N_1, \dots, N_k)}(U)$ is given by

$$\alpha = \sum_I \alpha_I dx^I$$

where $I : D(I) \subset [k] \rightarrow \mathbb{N}^+$, and for all $i \in D(I)$ we have $I(i) \in [N_i - 1]$, $\alpha_I \in C^\infty(U)$ and $dx^I = \prod_{i \in D(I)} d^{I(i)} x_i$.

We declare that $\deg(x_i) = 0$ and $\deg(d^k x_i) = k$ then $\Omega_{(N_1, \dots, N_k)}(\mathbb{R}^n)$ is a graded algebra with the product given by: for $\alpha, \beta \in \Omega_{(N_1, \dots, N_k)}(\mathbb{R}^n)$, $\alpha = \sum_I \alpha_I dx^I$ and $\beta = \sum_J \beta_J dx^J$

$$\alpha\beta = \sum_I (\alpha\beta)_I dx^I,$$

where

$$(\alpha\beta)_I = \sum_{K_1 \sqcup K_2 = D(I)} \text{sgn}(K_1, K_2, I) \alpha_{I|K_1} \beta_{I|K_2},$$

$$\text{and } \text{sgn}(K_1, K_2, I) = \prod_{\substack{i \in K_1, j \in K_2 \\ i > j}} (-1)^{I(i)I(j)}.$$

For $\alpha = \sum_I \alpha_I dx^I$ we define the form $d\alpha$ by

$$d\alpha = \sum_{1 \leq s \leq k} \frac{\partial \alpha_I}{\partial x_s} dx_s \wedge dx^I + \sum_{s \in D(I)} (-1)^{\sum_{t < s} I(t)} \alpha_I dx^{I+\delta_s}$$

where $\delta_s : [k] \rightarrow \mathbb{N}$ is given by $e_s(j) = \begin{cases} 1 & j = s \\ 0 & j \neq s. \end{cases}$

We have the following

Theorem 17 $\Omega_{(N_1, \dots, N_k)}(\mathbb{R}^n)$ is a $(N_1 + \dots + N_k - k)$ -dga.

proof Since $\Omega_{(N_1, \dots, N_k)}(x_1, \dots, x_k) = \Omega_{N_1}(x_1) \hat{\otimes} \dots \hat{\otimes} \Omega_{N_k}(x_k)$, we just need to consider the one variable case $\Omega_N(x) = \mathbb{R}[x, dx, \dots, d^{N-1}x] / \langle d^i x d^j x \rangle$, for $1 \leq i \leq N-1$. For $\alpha = \sum_{i=0}^{N-1} f_i(x) d^i(x)$ we have

$$d^k \alpha = \frac{\partial f_0}{\partial x} d^k x + \sum_{i=1}^{N-k-1} f_i(x) d^{i+k} x.$$

Taking $k = N$ we see that $\Omega_N(x)$ is a N -complex, the Leibniz rule and associativity of the product are easy to check, thus $\Omega_N(x)$ is a N -dga. By [AD, Theorem 8] $\Omega_{(N_1, \dots, N_k)}(\mathbb{R}^n)$ is a $(N_1 + \dots + N_k - k)$ -dga. $\blacklozenge$

We shall use the notation $\Omega_N(\mathbb{R}^k) := \Omega_{N, \dots, N}(\mathbb{R}^k)$.

Let us recall the definition of affine varieties.

Definition 18 M is an affine variety if there is an open covering $\Lambda = \{U_i\}$ of M and diffeomorphisms $\varphi_i : U_i \rightarrow \mathbb{R}^n$, with $\varphi_i(U_i)$ open, such that

$$\varphi_j \circ \varphi_i^{-1} : \varphi_i(U_i \cap U_j) \longrightarrow \varphi_j(U_i \cap U_j)$$

satisfy $\varphi_j \circ \varphi_i^{-1}(x) = A_{ij}x + b_{ij}$, $A_{ij} \in Gl_n(\mathbb{R})$ and $b_{ij} \in \mathbb{R}^n$.

$f : \mathbb{R}^k \rightarrow \mathbb{R}^k$ is an affine map, i.e., $f(x) = Ax + b$ for $A \in Gl_k(\mathbb{R}^k)$, $b \in \mathbb{R}^k$. The map given by

$$f^*(x_i) = f^i \quad \text{and} \quad f^*(d^j x_i) = \frac{\partial f^i}{\partial x_j} d^k x_j,$$

can be extended to a linear map $f : \Omega_N(\mathbb{R}^k) \rightarrow \Omega_N(\mathbb{R}^k)$ such that 1) $df^* = f^*d$ and 2) $f^*(\alpha \wedge \beta) = f^*(\alpha) \wedge f^*(\beta)$, for all $\alpha, \beta \in \Omega_N(\mathbb{R}^k)$.

Let M be an affine manifold. We define the algebra $\Omega_N(M)$ of differential forms of depth N on M as follows, $\Omega_N(M)$ consist of tuples $\alpha = (\alpha_U)_{U \in \Lambda}$, where $\alpha_U \in \Omega_N(\varphi_U(U))$, satisfying the following compatibility condition

$$(\varphi_V \circ \varphi_U^{-1})^*(\alpha_V|_{\varphi_V(U \cap V)}) = \alpha_U|_{\varphi_U(U \cap V)}.$$

For $U, V \in \Lambda$ such that $U \cap V \neq \emptyset$. Our final result is the following

Theorem 19 *Let M an affine manifold with $\dim(M) = m$. $\Omega_N(M)$ is a $(m-1)N$ -dga and $\mathbf{H}_N(M)$ is a $((m-1)N-1)$ -dga.*

References

- [A] V. Abramov, *On realizations of exterior calculus with $dN=0$* , Czechoslovak Journal of Physics, 48 (1998) 1265-1272.
- [AB] V. Abramov, N. Bazunova, *Exterior calculus with $d^3 = 0$ on a free associative algebra and reduced quantum plane*, Proceedings of XIV Max Born Symposium, New Symmetries and Integrable Models, Karpacz, Poland, 21-24.09.1999, World Scientific, Singapore-New Jersey-London-Hong Kong, 2000, pp. 3-7.
- [AK] V. Abramov, R. Kerner, *On certain realizations of the q -deformed exterior differential calculus*, Reports on Math. Phys., 43 (1999) 179-194
- [AD] M. Angel, R. Díaz, *N -differential graded algebras*, Preprint math.DG/0504398.
- [BGV] N. Berline, E. Getzler, M. Vergne, *Heat Kernels and Dirac Operators*, Grundlehren 298, Springer-Verlag, Berlin-Heidelberg-New York, 1992.
- [F] T. Frankel, *The geometry of physics: an introduction*, Cambridge University Press, 1997.
- [DV1] M. Dubois-Violette, *Generalized differential spaces with $d^N = 0$ and the q -differential calculus*, Czech J. Phys. 46 (1996) 1227- 1233.

- [DV2] M. Dubois-Violette, *Generalized homologies for $d^N = 0$ and graded q -differential algebras*, Contemporary Mathematics 219, American Mathematical Society 1998, M. Henneaux, J. Krasilshchik, A. Vinogradov, Eds., p. 69-79.
- [DV3] M. Dubois-Violette, *Lectures on differentials, generalized differentials and some examples related to theoretical physics*, Contemporary Mathematics 294, American Mathematical Society 2002, R. Coquereaux, A. Garcia, R. Trinchero Eds, p. 59-94.
- [DVK] M. Dubois-Violette, R. Kerner, *Universal q -differential calculus and q -analog of homological algebra*, Acta Math. Univ. Comenianae Vol. LXV, 2 (1996), 175-188.
- [K] M.M. Kapranov, *On the q -analog of homological algebra*, Preprint q-alg/9611005.
- [KW] C. Kassel, M. Wambst, *Algèbre homologique des N -complexes et homologie de Hochschild aux racines de l'unité*, Publ. Res. Inst. Math. Sci. Kyoto University 34 (1998), n 2, 91-114.
- [KN] R. Kerner, B. Niemeyer, *Covariant q -differential calculus and its deformations at $q^N = 1$* , Lett. in Math. Phys., 45, 161-176, (1998).
- [M] W. Mayer, *A new homology theory I, II*, Annals of Math. 43 (1942) 370-380 and 594-605.
- [S] A. Sitarz; *On the tensor product construction for q -differential algebras*, Lett. Math. Phys. 44 1998.

Mauricio Angel. Universidad Central de Venezuela (UCV). `mangel@euler.ciens.ucv.ve`
 Rafael Díaz. Universidad Central de Venezuela (UCV). `rdiaz@euler.ciens.ucv.ve`